

Monthly and Interannual Variability of the Eastern Pacific Warm Pool

Luis Zamudio¹ , Pat Hogan², Joe Metzger², & Jorge Zavala³

¹Center for Ocean-Atmospheric Prediction Studies, Florida State University

²Naval Research Laboratory, Stennis Space Center, Mississippi

³Centro de Ciencias de la Atmósfera, Universidad Nacional Autónoma de México

**HYCOM NOPP GODAE meeting, (Nov. 7-9, 2006)
COAPS, Florida State University, FL.**

Report Documentation Page			Form Approved OMB No. 0704-0188		
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE NOV 2006		2. REPORT TYPE		3. DATES COVERED 00-00-2006 to 00-00-2006	
4. TITLE AND SUBTITLE Monthly and Interannual Variability of the Eastern Pacific Warm Pool			5a. CONTRACT NUMBER		
			5b. GRANT NUMBER		
			5c. PROGRAM ELEMENT NUMBER		
6. AUTHOR(S)			5d. PROJECT NUMBER		
			5e. TASK NUMBER		
			5f. WORK UNIT NUMBER		
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Naval Research Laboratory,Stennis Space Center,MS,39529			8. PERFORMING ORGANIZATION REPORT NUMBER		
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)			10. SPONSOR/MONITOR'S ACRONYM(S)		
			11. SPONSOR/MONITOR'S REPORT NUMBER(S)		
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 16	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

Results indicate:


- The Eastern Pacific Warm Pool (EPWP) strengthens and weakens, but remains throughout the year.
- The monthly variability of the EPWP is forced by the surface heat fluxes.
- The interannual variability of the EPWP is modulated by the warm water advected poleward by interannual coastally trapped waves.

Tools


- 25 year non-assimilative simulation (1979-2003) of a Pacific configuration of HYCOM.
- 13 year (1993-2006) of Sea Surface Temperature data from the Modular Ocean Data Assimilation System (MODAS).

1/12° Pacific HYCOM Basin-scale Temperature


SST Climatological Mean (1979-2003) for May




SST Climatological Mean (1979-2003) for May




Tropical Cyclone Frequency





The eastern Tropical Pacific area of cyclone formation accounts for ~17% of the global total of tropical storm development (Amador et al., 2006).






- Why does the EPWP reach its maximum during May ?
- Why does the EPWP weaken during mid summer ?



SST > 28.5°C

- EPWP's May maximum of ~4,000,000 km²
- EPWP's January minimum of ~30,000 km²
- EPWP's July mid-summer minimum of ~2,300,000 km²
- EPWP gains heat through the surface from February to October
- EPWP loses heat from November to January.

SST > 28.5°C




MODAS is blue


HYCOM is red

- The monthly variability of the EPWP is forced by the surface heat fluxes.
- If the surface heat fluxes are the main forcing of the EPWP, then the surface heat fluxes should include an interannual variability, since the EPWP includes interannual variability in extension and strength.


Surface heat fluxes




Surface heat fluxes (no interannual variability)



Thermocline depth (interannual variability)




SST > 28.5°C (interannual variability)



Time

- Sea surface height anomaly time series from 1/16° Pacific NLOM, first along the equator (starting in the western Pacific and propagating eastward until arrival at the Americas West Coast), and second along the coast to the 20°N.




Summary

- The EPWP includes maximums and minimums of $\sim 4,000,000 \text{ km}^2$, and $\sim 30,000 \text{ km}^2$ during May and January, respectively and a mid-summer relative minimum of $\sim 2,300,000 \text{ km}^2$ during July. Those maximums and minimums are partially explained by the corresponding one month lagged maximums and minimums in the Surface heat fluxes.
- The extension of the EPWP has a strong interannual variability increasing (decreasing) during El Niño (La Niña) years. That is due to the interannual variability in the generation of Equatorial Pacific Kelvin waves.

Research in Progress

HYCOM-GLB-053 SSH and Currents for 2004-154-00



HYCOM-GLB-053 SSH and Currents for 2004-214-00

